

<https://helda.helsinki.fi>

Correction to: Predicting and clustering plant CLE genes with a new method developed specifically for short amino acid sequences

Zhang, Zhe

BioMed Central

2020-12-18

BMC Genomics. 2020 Dec 18;21(1):895

<http://hdl.handle.net/10138/323529>

Downloaded from Helda, University of Helsinki institutional repository.

This is an electronic reprint of the original article.

This reprint may differ from the original in pagination and typographic detail.

Please cite the original version.

CORRECTION

Open Access



Correction to: Predicting and clustering plant *CLE* genes with a new method developed specifically for short amino acid sequences

Zhe Zhang^{1,2}, Lei Liu^{1,2}, Melis Kucukoglu^{3,4}, Dongdong Tian^{1,2}, Robert M. Larkin^{1,2}, Xueping Shi^{1,2*} and Bo Zheng^{1,2*}

Correction to: BMC Genomics 21, 709 (2020)
<https://doi.org/10.1186/s12864-020-07114-8>

Following the publication of the original article [1], it was reported that there was an error in Fig. 7 whereby the yellow triangles showing the cleavage sites were not present. The corrected Fig. 7 is included in this Correction article, and the original article has been corrected.

Author details

¹Key Laboratory of Horticultural Plant Biology of Ministry of Education, Huazhong Agricultural University, Wuhan 430070, China. ²College of Horticulture and Forestry Sciences, Huazhong Agricultural University, Wuhan 430070, China. ³Institute of Biotechnology, Helsinki Institute of Life Science (HILIFE), University of Helsinki, 00014 Helsinki, Finland. ⁴Viikki Plant Science Centre, University of Helsinki, 00014 Helsinki, Finland.

Published online: 18 December 2020

Reference

1. Zhang Z, Liu L, Kucukoglu M, et al. Predicting and clustering plant *CLE* genes with a new method developed specifically for short amino acid sequences. *BMC Genomics*. 2020;21:709. <https://doi.org/10.1186/s12864-020-07114-8>.

The original article can be found online at <https://doi.org/10.1186/s12864-020-07114-8>.

* Correspondence: xpshi@mail.hzau.edu.cn; bozheng@mail.hzau.edu.cn

¹Key Laboratory of Horticultural Plant Biology of Ministry of Education, Huazhong Agricultural University, Wuhan 430070, China

Full list of author information is available at the end of the article



© The Author(s). 2020 **Open Access** This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>. The Creative Commons Public Domain Dedication waiver (<http://creativecommons.org/publicdomain/zero/1.0/>) applies to the data made available in this article, unless otherwise stated in a credit line to the data.




Fig. 7 Clustering analysis of IDA-like CLE motifs and Arabidopsis IDA/IDL motifs. **a** Clustering of IDA-like CLE motifs and Arabidopsis IDA/IDL, PIP/PIPL and CLV3 motifs. The heat map indicates the Euclidean distance of each pair of motifs. Red represents short distances. Blue represents long distances. A shorter Euclidean distance implies a higher similarity. **b** Protein domain schematic diagram of *Arabidopsis* IDA and two “PVPP-type” IDA-like CLE candidates. Protein domains were predicted using SMART. Blue box: RLK5-binding domain; red-brown box: low complexity domain; pale-brown triangle: location of the cleavage site of the signal peptide for the secretory pathway; black underline: IDA or IDA-like motif